

# Generation of Energetic Particles in Intense Laser Matter Interaction

B. Ramakrishna, M. Tayyab, S. Bagchi, T. Mandal,

J. A. Chakera, P. A. Naik, and P. D. Gupta

*Laser Plasma Division, Raja Ramanna Centre for Advanced Technology,  
Indore 452013, India*

\*E-mail: ramakrishnab@rrcat.gov.in

The acceleration of high energy ion beams (up to several tens of MeV per nucleon) following the interaction of an ultra-short ( $t < 50$  fs), intense ( $I\lambda^2 > 10^{18}$  W.cm<sup>-2</sup>.μm<sup>-2</sup>) laser pulse with solid targets, is one of the burgeoning fields of research in the last few years. Mechanisms leading to forward-accelerated, high quality ion beams, operating at currently accessible laser intensities (up to  $10^{21}$  W/cm<sup>2</sup>) in laser-matter interactions, are mainly associated with large electric fields set up at the target rear interface by the laser-accelerated electrons leaving the target. The emitted ion pulses, and in particular, the proton pulses contain a large number of particles (up to  $10^{13}$ ) with energies in excess of several MeV, having a pulse duration  $\sim$  ps, and a source size of tens to hundreds of μm [1-2]. Conversion efficiencies (laser energy to proton energy) up to 7 percent have been reported [3]. However, the emitted proton beam has a large divergence, which restricts its applications in ion cancer therapy and inertial confinement fusion (ICF) [4-5].

In this paper, we present our recent experimental results on MeV ion generation by mildly relativistic ( $10^{18}$  W/cm<sup>-2</sup>) short-pulse (45 fs) laser interaction with foil targets of varying thicknesses, structured / uniform targets (e.g. nano structures on thin metallic foils, sandwich targets). Spectral modification / bunching, and divergence from structured targets will be discussed. Observation of energetic negative hydrogen ions from ultra-short laser pulse interaction with thin foil targets will also be discussed.

- [1] R. A. Snavely *et al.*, Phys. Rev. Lett. **85**, 2945 (2000).
- [2] L. Romagnani *et al.*, Phys. Rev. Lett. **95**, 195001 (2005).
- [3] J. Fuchs *et al.*, Nature Phys. **2**, 48 (2006).
- [4] A. Macchi, M. Borghesi, and M. Passoni, Rev. Mod. Phys. **85**, 751 (2013).
- [5] K. W. D. Ledingham, W. Galster, and R. Sauerbrey, Br. J. Radiol. **80**, 855 (2007).