

# Study of an Atmospheric Pressure Argon Microplasma Jet

J. Amorim<sup>1,2</sup>, M. A. Ridenti<sup>2</sup> and V. Guerra<sup>3</sup>

<sup>1</sup>*Departamento de Física, Instituto Tecnológico de Aeronáutica, 12228-900, S.J. Campos, Brazil*

<sup>2</sup>*Instituto de Física Gleb Wataghin, UNICAMP, 13083-859, Campinas, Brazil*

<sup>3</sup>*Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico, Universidade de Lisboa, 1049-001 Lisboa, Portugal*

A surface wave discharge (SWD) in argon at atmospheric pressure generated by a Surfatron device, such as those used in the treatment of biomass for the production of bioethanol [1,2], is experimentally studied by optical emission and mass spectrometries. The measurements are complemented with the results from a self-consistent model of these discharges.

The electron density  $n_e$ , electron temperature  $T_e$  and gas temperature  $T_g$  were measured as a function of the applied power and gas flow rate.  $T_e$  was estimated using a collisional-radiative model that takes as input four emission lines originating from  $4p$  states [1,3], while  $n_e$  was determined through the Stark broadening of the  $H_\beta$  line profile. The self-absorbing method was used to estimate the population of the metastable state  $Ar(1s_5)$  [2,4]. The gas temperature was obtained from the rotational spectra of OH. The axial profile of  $n_e$ , from the launcher to the end of the plasma column, presents a maximum value under certain conditions, in contrast with the typical  $n_e$  profiles of SWDs, which are usually monotonically decreasing. A correlation between  $n_e$  and the metastable state  $Ar(1s_5)$  was found, suggesting that stepwise ionization from metastable states may play an important role [1].

Employing mass-resolved ion-energy distribution measurements, it was possible to study the behavior of fourteen ionic species ( $Ar^+$ ,  $Ar_2^+$ ,  $ArH^+$ ,  $H_2O^+$ ,  $H^+(H_2O)$ ,  $H^+(H_2O)_2$ ,  $O^+$ ,  $O_2^+$ ,  $OH^+$ ,  $NO^+$ ,  $NO_2^+$ ,  $N^+$ ,  $N_2^+$ ,  $N_4^+$ ) of the plasma in diffuse and contracted modes [2]. The crucial role of three molecular ions ( $Ar_2^+$ ,  $ArH^+$ ,  $H_2O^+$ ) during the radial contraction and the relationship between power, gas flow and molecular ions production is pointed out. Ion energy distributions for  $Ar_2^+$ ,  $ArH^+$  and  $H_2O^+$  exhibit their maxima at high-energies, due to the low-energy threshold for three-body collision reactions.

A first modeling effort is done with the development of a self-consistent model of an SWD in Ar at atmospheric pressure. The model is based on the solution of the homogeneous electron Boltzmann equation, considering inelastic and superelastic collisions with the  $Ar(4s)$  states and electron-electron collisions, coupled with a system of rate balance equations describing the creation and destruction of the most important heavy particles, namely ground-state  $Ar(^1S_0)$  atoms, the four  $Ar(4s)$  states,  $Ar(4p)$  states, treated here as a single effective level, the molecular dimer  $Ar_2^*$ , as well as  $Ar^+$  and  $Ar_2^+$  ions. The simulations and experimental results are compared and discussed, providing physical insight into the basic mechanisms and phenomena ruling the discharge.

**Acknowledgement:** VG was partially supported by FCT (Pest-OE/SADG/LA0010/2013)

- [1] M. A. Ridenti, J. A. Souza-Corrêa and J. Amorim, *J. Phys. D: Appl. Phys.* **47**, 045204 (2014).
- [2] M. A. Ridenti, N. Syrou and J. Amorim, *Chem. Phys. Lett.* **435**, 1 (2014).
- [3] D. Mariotti, Y. Shimiz, T. Sasaki and N. Koshizaki, *Appl. Phys. Lett.* **89**, 201502 (2006).
- [4] J. Jolly and M. Touzeau, *J. Quant. Spectrosc. Radiat. Transfer* **15**, 863 (1975).