

# Recombination of $\text{H}_3^+$ with electrons in afterglow plasma in hydrogen

Juraj Glosík, Petr Dohnal, Peter Rubovič, Ábel Kálosi, Michal Hejduk and Radek Plašil

*Charles University in Prague, Faculty of Mathematics and Physics, Prague, Czech Republic*

Recombination of  $\text{H}_3^+$  ions with electrons in stationary afterglow plasma in hydrogen at 240–340 K was studied using near-infrared (NIR) absorption spectroscopy (SA-CRDS). The plasma is formed in a pulsed microwave discharge in a mixture of He-Ar- $\text{H}_2$  and the densities of three rotational states of the ground vibrational state of  $\text{H}_3^+$  during the afterglow are measured using the transitions  $3v_2^1(2,0) \leftarrow 0v_2^0(1,0)$  and  $3v_2^1(4,3) \leftarrow 0v_2^0(3,3)$  for ortho- $\text{H}_3^+$  and  $3v_2^1(2,1) \leftarrow 0v_2^0(1,1)$  for para- $\text{H}_3^+$ . Details and notation are discussed in refs. [1,2]. The decay of the plasma is then analysed to obtain an “effective” recombination coefficient  $\alpha_{\text{eff}}$ . From dependence of the measured  $\alpha_{\text{eff}}$  on hydrogen density  $[\text{H}_2]$  binary rate coefficient ( $\alpha_{\text{bin}}$ ) and ternary  $\text{H}_2$ -assisted recombination rate coefficient ( $K_{\text{H}_2}$ ) are obtained. We found that the ternary process is by five orders of magnitude more effective than process described by classical theory of Bates and Khare [3] for ternary recombination. The value of the binary recombination rate coefficient is in agreement with generally accepted value [4,5,6].

This work was supported by GACR P209-12-0233, GACR 14-14649P, GAUK 642214, and GAUK 659112.

- [1] Hlavenka P., Plasil R., Bano G., Korolov I., Gerlich D., Ramanlal J., Tennyson J., Glosik J., *Int. J. Mass Spectrom.* **255**, 170–176 (2006)
- [2] Lindsay C. M., McCall B. J., *J. Mol. Spectrosc.* **210**, 60 (2001)
- [3] D. Bates and S. Khare, *Proc. Phys. Soc. Lond.* **85**, 231 (1965)
- [4] Johnsen, R.; Guberman, S. L. Dissociative Recombination of  $\text{H}_3^+$  Ions with Electrons: Theory and Experiment. *Adv. At., Mol., Opt. Phys.* **59**, 75–128 (2010)
- [5] Dohnal, P., Hejduk, M., Varju, J., *et al.* Binary and ternary recombination of para- $\text{H}_3^+$  and ortho- $\text{H}_3^+$  with electrons: State selective study at 77–200 K. *Journal of Chemical Physics* **136**, 244304 (2012)
- [6] Rubovič, P., Dohnal, P., Hejduk, M., *et al.* Binary recombination of  $\text{H}_3^+$  and  $\text{D}_3^+$  Ions with Electrons in Plasma at 50–230 K. *Journal of Physical Chemistry A* **117**, 86 (2013)